

Semi-stable fibrations of generic p -rank 0

Junmyeong Jang

email : jang3@math.purdue.edu

Mathematics Subject Classification : 11G25, 14J20

1 Introduction

Let k be an algebraically closed field and $\pi : X \rightarrow C$ be a semi-stable fibration of a connected proper smooth surface to a connected proper smooth curve over k . If the base field k is a subfield of $\mathbb{C}$, the field of complex numbers, the following semi-positivity theorem holds.

Theorem.(Semi-Positivity Theorem, Xiao) If $\pi : X \rightarrow C$ is a fibration of a proper smooth surface to a proper smooth curve over $\mathbb{C}$, then all the quotient bundles of $\pi_*\omega_{X/C}$ are of non-negative degree. [10],p.1

In general the semi-positivity theorem is not valid over a field of positive characteristic. In [7], Moret-Bailly constructed a semi-stable fibration $\pi_M : X_M \rightarrow \mathbb{P}^1$ of fiber genus 2 such that $R^1\pi_{M*}\mathcal{O}_{X_M} = \mathcal{O}(1) \oplus \mathcal{O}(-p)$ where p is the characteristic of the base field. In the previous work,[6] we have proved that for a semi-stable fibration $\pi : X \rightarrow C$, if the generic fiber is ordinary, then the semi-positivity theorem holds. Precisely, when the generic fiber of π is ordinary, all the Harder-Narasimhan slopes of $R^1\pi_*\mathcal{O}_X$ are non-positive. For Moret-Bailly's example, the p -rank of the generic fiber is 0. In particular, every special fiber of π_M is a supersingular smooth curve of genus 2 or a union of two supersingular elliptic curves which intersect at a point transversally. In this paper we prove the following theorem which generalizes the failure of the semi-positivity theorem for Moret-Bailly's example.

Theorem 1. Let $\pi : X \rightarrow C$ be a non-isotrivial semi-stable fibration of proper smooth surface to a proper smooth curve over a field of positive characteristic. If the generic p -rank of π is 0, then $F_C^{n*}R^1\pi_*\mathcal{O}_X$ has a positive Harder-Narasimhan slope for a sufficiently large $n \in \mathbb{N}$. In particular, if the genus of C is 0 or 1, $R^1\pi_*\mathcal{O}_X$ has a positive Harder-Narasimhan slope.

As an application of the theorem, we obtain a result on a distribution of p -ranks of reductions of a certain non-closed point in the moduli space of curves over a number field.

Corollary 2.8. Suppose that $\pi : X \rightarrow C$ is a non-isotrivial semi-stable fibration of base genus 0 or 1 defined over a number field F , and that $U \subseteq C$ is the smooth locus of π . π defines a non-constant morphism $f : U \rightarrow \mathcal{M}_{g,F}$. Let P be the image of the generic point of U under f . Then the reduction P_v is not contained in the p -rank 0 strata for almost all v .

This result can be considered as a variation of Serre's ordinary reduction conjecture. It is a weak statement since it is only about some non-closed points in the moduli space and 0 p -rank. But it is a somewhat interesting phenomenon that the semi-positivity theorem, which is concerned with a coherent module on a fiber of characteristic 0, encodes an information of p -ranks of the reductions.

2 Proof of Theorem 1.

We follow the terminology of [6]. Let us recall the definition of a semi-stable curve. Let k be an algebraically closed field and C be a projective curve over k .

Definition 2.1. C is (semi-)stable if

1. It is connected and reduced.
2. All the singular points are normal crossing.
3. An irreducible component, which is isomorphic to $\mathbb{P}^1$, meets other components in at least 3 (resp. 2) points.

For an arbitrary base scheme, we define a (semi-)stable curve as follows.

Definition 2.2. A proper flat morphism of relative dimension 1 of schemes $\pi : X \rightarrow S$ is a (semi-)stable curve if every geometric fiber of π is a (semi-)stable curve in the sense of definition 2.1.

In this paper, we assume $\pi : X \rightarrow C$ is a generically smooth semi-stable fibration of a proper smooth surface to a proper smooth curve over a field k unless it is stated otherwise.

Definition 2.3. For a generically smooth semi-stable fibration $\pi : X \rightarrow C$ defined over a field of positive characteristic, the generic p -rank of π is the p -rank of a geometric generic fiber of π .

2.1 Self duality of $B\omega^1$

Let k be a perfect field of positive characteristic and $\pi : X \rightarrow C$ be a generically smooth semi-stable curve. Let $\omega_{X/C}^1$ be the relative dualizing line bundle for π . There is the canonical inclusion $i : \Omega_{X/C}^1 \hookrightarrow \omega_{X/C}^1$. At a relative smooth point for π , i is an

isomorphic. On the other hand, at a relative singular point, where étale locally π is given by

$$\mathrm{Spec} k[x, y, t]/(xy - t) \rightarrow \mathrm{Spec} k[t],$$

$\omega_{X/C}^1$ is a free module of rank 1 generated by $dx/x = -dy/y$ and $\Omega_{X/C}^1$ is a submodule of $\omega_{X/C}^1$ generated by dx and dy via i . Composing with the inclusion i , we have the differential morphism $d : \mathcal{O}_X \rightarrow \omega_{X/C}^1$. When $F_{X/C}$ is the relative Frobenius morphism for π , π

$$\begin{array}{ccccc} X & \xrightarrow{F_{X/C}} & X^p & \xrightarrow{\quad} & X \\ & \searrow \nearrow & \downarrow \pi^p & & \downarrow \pi \\ & & C & \xrightarrow{F_C} & C, \end{array}$$

$d : F_{X/C*}\mathcal{O}_X \rightarrow F_{X/C*}\omega_{X/C}^1$ is $\mathcal{O}_{X^p}$ -linear. The kernel of d is the image of $F_{X/C}^* : \mathcal{O}_{X^p} \hookrightarrow F_{X/C*}\mathcal{O}_X$ and the image of d is denoted by $B^1\omega_{X/C}$ or $B^1\omega$. $B\omega^1$ is flat over $\mathcal{O}_C$. Let $U \hookrightarrow X$ be the smooth locus for π . The usual Cartier isomorphism

$$C : \Omega_{U/C}^1/B^1\Omega_{U/C} \rightarrow \Omega_{U^p/C}$$

is extended to an isomorphism

$$C : \omega_{X/C}^1/B^1\omega \rightarrow \omega_{X^p/C}^1. [5], \text{p.381}$$

Using this Cartier morphism, we have an $\mathcal{O}_{X^p}$ -linear paring

$$F_{X/C*}\mathcal{O}_X \otimes F_{X/C*}\omega_{X/C}^1 \rightarrow \omega_{X^p/C}^1, (\alpha, \omega) \mapsto C(\alpha\omega).$$

This pairing induces a pairing

$$(F_{X/C*}\mathcal{O}_X/\mathcal{O}_{X^p}) \otimes B^1\omega \rightarrow \omega_{X^p/C}^1.$$

On U , this pairing gives a perfect self duality. In particular, we have

$$B^1\Omega_{U/C} \simeq \underline{Hom}(B^1\Omega_{U/C}, \Omega_{U^p/C}^1).$$

Proposition 2.4. If X is a smooth surface over a perfect field k which admits a semi-stable fibration, $\pi : X \rightarrow C$, to a smooth curve C over k , then

$$B^1\omega_{X/C} \simeq \underline{Hom}(B^1\omega_{X/C}, \omega_{X^p/C}^1).$$

Proof. It's enough to check that the paring is perfect at the relative singular points in X . Let $x \in X$ be a relative singular point. Étale locally, we may assume $X = \mathrm{Spec} A$, $C = \mathrm{Spec} B$ where

$$A = k[x, y, t]/(xy - t) \simeq k[x, y], \quad B = k[t]$$

and π is the canonical morphism

$$k[t] \rightarrow k[x, y, t]/(xy - t).$$

Let $A^p = A \otimes_B (B, F_B)$. Then $A^p = k[X, Y, t]/(XY - t^p)$ and the relative Frobenius morphism is a k -algebra morphism $F_{A/B} : A^p \rightarrow A$ given by

$$X \mapsto x^p, Y \mapsto y^p \text{ and } t \mapsto xy.$$

We may regard A^p is a k -subalgebra of A generated by x^p, y^p, xy . As an A^p -module, $B^1\omega = A/A^p$ is generated by

$$x, x^2, \dots, x^{p-1}, y, \dots, y^{p-1}.$$

$B^1\omega$ is a torsion free A^p -module and the $\text{Frac}(A^p)$ -dimension of $B^1 \otimes_{A^p} \text{Frac}(A^p)$ is $p - 1$. Therefore there are only $p - 1$ obvious relations

$$t^{p-1}x = Xy^{p-1}, t^{p-2}x^2 = Xy^{p-2}, \dots, tx^{p-1} = Xy$$

among the generators and we have an A^p -module decomposition

$$B^1\omega = \bigoplus_{i=1}^{p-1} \langle x^i, y^{p-i} \rangle.$$

On the other hand, $\omega_{A/B}^1$ is a rank 1 free A -module generated by $dx/x = -dy/y$ and $\omega_{A^p/B}^1$ is a rank 1 free A^p -module generated by $dX/X = -dY/Y$. The Cartier morphism is the A^p -linear morphism satisfying

$$\begin{aligned} dx/x &\mapsto dX/X, \\ xdx/x &\mapsto 0, \\ &\vdots \\ x^{p-1}dx/x &\mapsto 0, \\ ydx/x &\mapsto 0, \\ &\vdots \\ y^{p-1}dx/x &\mapsto 0. \end{aligned}$$

In the above decomposition of $B^1\omega$, it's easy to see that the dual of $\langle x^i, y^{p-i} \rangle$ is $\langle x^{p-i}, y^i \rangle$ and that the pairing $B^1\omega \otimes B^1\omega \rightarrow \omega_{X^p/C}^1$ gives a perfect duality of the dual components of both sides. This proves the claim. $\square$

Corollary 2.5. Let k be a perfect field of positive characteristic. Let $\pi : X \rightarrow C$ be a semi-stable fibration of a proper smooth surface to a proper smooth curve over k . Let M and T be the free part and the torsion part of $R^1\pi_*B^1\omega_{X/C}$ respectively and $N = R^1\pi_*\mathcal{O}_X$. Then there exists a exact sequence of coherent modules on C

$$0 \rightarrow M^* \rightarrow F_C^*N \rightarrow N \rightarrow M \oplus T \rightarrow 0.$$

Proof. The exact sequence of coherent $\mathcal{O}_{X^p}$ -modules

$$0 \rightarrow \mathcal{O}_{X^p} \rightarrow F_{X/C*} \mathcal{O}_X \rightarrow B^1 \omega_{X/C} \rightarrow 0$$

gives a long exact sequence for the π_*^p functor

$$0 \rightarrow \mathcal{O}_C \cong \mathcal{O}_C \rightarrow \pi_*^p B^1 \omega_{X/C} \rightarrow R^1 \pi_*^p \mathcal{O}_{X^p} \rightarrow R^1 \pi_*^p F_{X/C*} \mathcal{O}_X \rightarrow R^1 \pi_*^p B^1 \omega_{X/C} \rightarrow 0.$$

Because the Frobenius morphism of C is finite flat, $R^1 \pi_*^p \mathcal{O}_{X^p} = F_C^* R^1 \mathcal{O}_X$. And the relative Frobenius morphism $F : X \rightarrow X^p$ is finite affine, so $R^1 \pi_*^p F_{X/C*} \mathcal{O}_X = R^1 \pi_* \mathcal{O}_X$. By proposition 2.4, $B^1 \omega_{X/C} \simeq \underline{\text{Hom}}(B^1 \omega_{X/C}, \omega_{X^p/C})$. Since π is relative 1-dimensional, by the relative duality theorem

$$\pi_*^p B^1 \omega_{X/C} = \underline{\text{Hom}}(R^1 \pi_*^p B^1 \omega_{X/C}, \mathcal{O}_C).$$

Therefore the claim follows. $\square$

2.2 Proof of the theorem

Theorem 1. Let $\pi : X \rightarrow C$ be a non-isotrivial semi-stable fibration of proper smooth surface to a proper smooth curve over a field of positive characteristic. If the generic p -rank of π is 0, then $F_C^{n*} R^1 \pi_* \mathcal{O}_X$ has a positive Harder-Narasimhan slope for a sufficiently large $n \in \mathbb{N}$. In particular, if the genus of C is 0 or 1, $R^1 \pi_* \mathcal{O}_X$ has a positive Harder-Narasimhan slope.

Proof. The n -iterative relative Frobenius morphism in the diagram

$$\begin{array}{ccccc} X & \xrightarrow{F_{X/C}^n} & X^{p^n} & \xrightarrow{\quad} & X \\ & \searrow & \downarrow \pi^{p^n} & & \downarrow \pi \\ & & C & \xrightarrow{F_C^n} & C \end{array}$$

is the composition of relative Frobenius morphisms

$$X \xrightarrow{F_{X/C}} X^p \xrightarrow{F_{X^p/C}} \dots \rightarrow X^{p^{n-1}} \xrightarrow{F_{X^{p^{n-1}}/C}} X^{p^n}.$$

$F_{X/C}^n$ gives an exact sequence of coherent $\mathcal{O}_{X^{p^n}}$ -modules

$$(*) \quad 0 \rightarrow \mathcal{O}_{X^{p^n}} \xrightarrow{F^{n*}} F_*^n \mathcal{O}_X \rightarrow E_n \rightarrow 0.$$

Here E_n is flat over $\mathcal{O}_C$ and $E_1 = B^1 \omega_{X/C}$. If we denote $N = R^1 \pi_* \mathcal{O}_X$, $R^1 \pi_*^{p^n} \mathcal{O}_{X^{p^n}} = F_C^{n*} N$. Let $\lambda_n : F_C^{n*} N \rightarrow N$ be the morphism induced by $F_{X/C}^{n*}$ in $(*)$. Because the relative Frobenius morphism commutes with a base change and the Frobenius morphism

of C is flat, λ_n is the composition of Frobenius pullbacks of $\lambda_1 : F_C^* N \rightarrow N$,

$$\lambda_n = \lambda_1 \circ \cdots \circ F_C^{(n-2)*} \lambda_1 \circ F_C^{(n-1)*} \lambda_1 : F_C^{n*} N \rightarrow F_C^{(n-1)*} N \rightarrow \cdots \rightarrow F_C^* N \rightarrow N.$$

On the other hand, the restriction of $(*)$ to a special fiber X_s is

$$0 \rightarrow \mathcal{O}_{X_s^{p^n}} \rightarrow F_{X_s/k(s)}^n \mathcal{O}_{X_s} \rightarrow E_{n,s} \rightarrow 0.$$

Furthermore we have the long exact sequence

$$\cdots \rightarrow H^1(\mathcal{O}_{X_s^{p^n}}) \rightarrow H^1(\mathcal{O}_{X_s}) \rightarrow H^1(E_{n,s}) \rightarrow 0.$$

Since we have assumed that the generic p -rank of π is 0, by the Grothendieck specialization theorem, for all $s \in C$, the p -rank of X_s is 0. It follows that there exists $n \in \mathbb{N}$, such that $H^1(X_s, \mathcal{O}_{X_s^{p^n}}) \rightarrow H^1(X_s, \mathcal{O}_{X_s})$ is the zero morphism for all $s \in C$ such that X_s is smooth. Hence $\dim H^0(X_s, E_{n,s}) = \dim H^1(X_s, E_{n,s}) = \text{gen } X_s = g$ for such s . But by the semi-continuity theorem, $\pi_*^{p^n} E_n$ and $R^1 \pi_*^{p^n} E_n$ are vector bundles of rank g . Because N is also a vector bundle of rank g on C , considering the exact sequence

$$0 \rightarrow \pi_*^{p^n} E_n \rightarrow F_C^{n*} N \xrightarrow{\lambda_n} N \rightarrow R^1 \pi_*^{p^n} E_n \rightarrow 0,$$

$\lambda_n = 0$. Now assume all the Harder-Narasimhan slopes of $F_C^i N$ are non-positive for all i . Let M be the free part of $R^1 \pi_*^p B^1 \omega_{X/C}$. Then $\pi_*^p B^1 \omega_{X/C} = M^* = \underline{\text{Hom}}_{\mathcal{O}_C}(M, \mathcal{O}_C)$. (Corollary 2.4) All the Harder-Narasimhan slopes of $F_C^{i*} M^*$ are non-positive since $F_C^{i*} M^*$ is a sub-bundle of $F_C^{i*} N$, so all the Harder-Narasimhan slopes of $F_C^{i*} M$ are non-negative. Let us consider an exact sequence

$$0 \rightarrow \text{Im } \lambda_{i-1} / \text{Im } \lambda_i \rightarrow N / \text{Im } \lambda_i \rightarrow N / \text{Im } \lambda_{i-1} \rightarrow 0. (i \geq 2)$$

We can also think of the free part of the above exact sequence

$$0 \rightarrow V_i' \rightarrow V_i \rightarrow V_i'' \rightarrow 0.$$

Here V_i is the free part of $N / \text{Im } \lambda_i$ and V_i'' is the free part of $N / \text{Im } \lambda_{i-1}$. V_i' is the saturation of the free part of $\text{Im } \lambda_{i-1} / \text{Im } \lambda_i$ in V_i . We can see $V_2'' = M$ is of non-negative degree by the assumption. On the other hand, since V_i' is a saturation of a quotient bundle of $F^{(i-1)*} M$, it is also of non-negative degree. Therefore, by induction V_i is of non-negative degree. Since $\lambda_n = 0$ for a sufficiently large n , the degree of $V_n = N$ is non-negative. But since π is non-isotrivial, $\deg N$ is strictly negative. [9], p.173 It is contradiction, so $F^{i*} N$ has a positive Harder-Narasimhan slope for some i . If the genus of the base C is 0 or 1, the Frobenius pull back F_C^* preserves the semi-stability of vector bundles, so $R^1 \pi_* \mathcal{O}_X$ has a positive Harder-Narasimhan slope. $\square$

Remark 2.6. The failure of the semi-positivity theorem for Moret-Bailly's example in the introduction is a special case of Theorem 2. Using Theorem 2, we may construct a lot of counterexamples of the semi-positivity theorem. Assume k is algebraically

closed of positive characteristic. In $\mathcal{M}_{g,k}$, the moduli space of smooth proper curves of genus g over k , the p -rank 0 strata is a closed subscheme which is purely $2g - 3$ dimensional.[3],p.120 Let P be a 1 dimensional point in the p -rank 0 strata. By the semi-stable reduction theorem,[1],p.3 there is a semi-stable fibration $\pi : X \rightarrow C$ such that the morphism $C \rightarrow \mathcal{M}_{g,k}$ induced by π sends the generic point of C to P . Since the generic p -rank of π is 0, for a suitable Frobenius base change $\pi^{p^n} : X^{p^n} \rightarrow C$, $R^1\pi_* \mathcal{O}_{X^{p^n}}$ has a positive slope by Theorem 1. X^{p^n} may contain isolated singularities. But the composition of π^{p^n} and the desingularization $X^{(n)} \rightarrow X^{p^n}$, $\pi^{(n)} : X^{(n)} \rightarrow C$ is a semi-stable fibration and $R^1\pi_*^{(n)} \mathcal{O}_{X^{(n)}} = R^1\pi_*^{p^n} \mathcal{O}_{X^{p^n}}$. Hence $\pi^{(n)}$ is a counterexample to the semi-positivity theorem.

Corollary 2.7. Let F be a number field and suppose a semi-stable fibration $\pi : X \rightarrow C$ is defined over F . There is an integral model of π , $\pi_A : X_A \rightarrow C_A$ defined over $\text{Spec } A$, an affine open set of $\text{Spec } \mathcal{O}_F$. Let $\pi_v : X_v \rightarrow C_v$ be the reduction of π_A at a place $v \in \text{Spec } A$. If the genus of C is 0 or 1, then the generic p -rank of π_v is not 0 for all but finitely many places v .

Proof. Since the harder-Narasimhan filtration of $R^1\pi_{A*} \mathcal{O}_{X_A}$ on the generic fiber of $\mathbb{C}_A \rightarrow \text{Spec } A$ extends to a non-empty open set of $\text{Spec } A$, $R^1\pi_{v*} \mathcal{O}_{X_v}$ has no positive Harder-Narasimhan slope for almost all $v \in \text{Spec } A$. Therefore the generic p -rank of π_v is not 0 by Theorem 1. $\square$

Let $\mathcal{M}_{g,\mathcal{O}_F}$ be the moduli space of proper smooth curves over $\text{Spec } \mathcal{O}_F$. $\mathcal{M}_{g,F}$, the moduli space over F , is the generic fiber of $\mathcal{M}_{g,\mathcal{O}_F} \rightarrow \text{Spec } \mathcal{O}_F$. When P is a geometrically irreducible point of $\mathcal{M}_{g,F}$, the closure of P in $\mathcal{M}_{g,\mathcal{O}_F}$ has a geometrically irreducible reduction at almost all places v . Let us denote the generic point of the reduction at v by P_v . P_v is contained in a p -rank strata in $\mathcal{O}_{g,\bar{k}_v}$, the moduli space over the residue field at v . We may ask the distribution of the p -ranks of P_v . Serre's ordinary reduction conjecture is a problem for closed points in the moduli space. In the language of the moduli space, Corollary 2.7 can be stated as follow.

Corollary 2.8. Suppose that $\pi : X \rightarrow C$ is a non-isotrivial semi-stable fibration of base genus 0 or 1 defined over a number field F , and that $U \subseteq C$ is the smooth locus of π . π defines a non-constant morphism $f : U \rightarrow \mathcal{M}_{g,F}$. Let P be the image of the generic point of U under f . Then the reduction P_v is not contained in the p -rank 0 strata for almost all v .

Remark 2.9. If the fiber genus of $\pi : X \rightarrow C$ is 2, Corollary 2.7 holds for arbitrary base C . Ekedahl showed that if $\pi : X \rightarrow C$ is a generically super-singular semi-stable fibration of fiber genus 2, then there exist a finite étale cover $f : D \rightarrow C$ and a morphism $g : D \rightarrow \mathbb{P}^1$ such that $\pi_f : X \times_C D \rightarrow D$ is isomorphic to $\pi_{M,g} : X_M \times_{\mathbb{P}^1} D \rightarrow D$ where $\pi_M : X_M \rightarrow \mathbb{P}^1$ is Moret-Bailly's fibration.[2],p.173 Since a pullback by a finite separable morphism of curves preserves the semi-stability of vector bundles, $R^1\pi_* \mathcal{O}_X$ has a positive Harder-Narasimhan slope. Considering the construction of a semi-stable fibration from a 1-dimensional point in the moduli space(Remark 2.6) and the Grothendieck specialization theorem, Corollary 2.8 holds for an arbitrary non-closed point in $\mathcal{M}_{2,F}$.

It is natural to expect that Corollary 2.8 holds for an arbitrary non closed point in $\mathcal{M}_{g,F}$ for any g , or equivalently that Corollary 2.7 holds for arbitrary base curve C . If the result of Proposition 2.12 in [6] is valid over a field of characteristic 0, i.e. the slope 0 part of $R^1\pi_*\mathcal{O}_X$ is potentially trivial, this expectation is valid. Indeed, in the situation of Corollary 2.7 without the assumption of the base genus, if the slope 0 part, $(R^1\pi_*\mathcal{O}_X)_0$ is potentially trivial, $(R^1\pi_v\mathcal{O}_{X_v})_0$ is strongly semi-stable for almost all $v \in \text{Spec } A$. On the other hand, by [8], p.660, the negative slope part, $F_v^{n*}(R^1\pi_*\mathcal{O}_{X_v})_-$ has only negative Harder-Narasimhan slopes for any n if the residue characteristic of v is sufficiently large. Therefore the claim follows.

Question 2.10. For an arbitrary geometrically irreducible non-closed point P in $\mathcal{M}_{g,F}$, is the p -rank of the reduction P_v nonzero for almost all $v \in \text{Spec } \mathcal{O}_F$?

References

- [1] Deschamps, M. Réduction Semi-Stable, *Asterisque* 86, 1981, 1-34.
- [2] Ekedahl, T. On Supersingular Curves and Abelian Varieties, *Math.Scand*, 60, 1987, 151-178.
- [3] Faber, C. and Van Der Geer, G. Complete subvarieties of moduli spaces and the Prym map, *J.Reine Angew. Math.* 573, 2004, 117-137.
- [4] Fujita, T. On Kähler fiber spaces over curves, *J.Math.Soc.Japan*, 30, no.4, 1978, 779-794.
- [5] Illusie, L. Ordinarité des intersections complètes générales, *The Grothendieck Festschrift vol.II*, *Progress in Mathematics* 87, 1990, 375-405.
- [6] Jang, J. Generic ordinarity for semi-stable fibrations, preprint, 2008.
- [7] Moret-Bailly, L. Familles de Courbes et de Variétés Abéliennes sur $\mathbb{P}^1$, *Asterisque* 86, 1981, 125-140.
- [8] Shepherd-Barron, N. Semi-stability and Reduction mod p , *Topology* 37, 1998, 659-664.
- [9] Szpiro, L. Sur Le Théorème de rigidité de Parsin et Arakelov, *Asterisque* 64, 1979, 169-202.
- [10] Xiao, G. Surfaces fibrees en courbes de genre deux, *Lecture Notes in Mathematics* 1137, 1980.